arXiv:1403.6640v1 [math.AG] 26 Mar 2014

ON THE GROUP OF AUTOMORPHISMS OF A QUASI-AFFINE VARIETY

ZBIGNIEW JELONEK

Abstract. Let $\mathbb{K}$ be an algebraically closed field of characteristic zero. We show that if the automorphisms group of a quasi-affine variety X is infinite, then X is uniruled.

1. Introduction.

Automorphism groups of an open varieties have always attracted a lot of attention, but the nature of this groups is still not well-known. For example the group of automorphisms of $\mathbb{K}^n$ is understood only in the case $n = 2$ (and $n = 1$, of course). Let Y be a an open variety. It is natural to ask when the group of automorphisms of Y is finite. We gave a partial answer to this questions in our papers [Jel1], [Jel2], [Jel3] and [Jel4]. In [Iit2] Iitaka proved that $Aut(Y)$ is finite if Y has a maximal logarithmic Kodaira dimension. Here we focus on the group of automorphisms of an affine or more generally quasi-affine variety over an algebraically closed field of characteristic zero. Let us recall that a quasi-affine variety is an open subvariety of some affine variety. We prove the following:

Theorem 1.1. *Let X be a quasi-affine (in particular affine) variety over an algebraically closed field of characteristic zero. If $Aut(X)$ is infinite, then X is uniruled, i.e., X is covered by rational curves.*

This generalizes our old results from [Jel3] and [Jel4]. Our proof uses in a significant way a recent progress in a minimal model program (see [Bir], [BCHK], [P-S]) and is based on our old ideas from [Jel1], [Jel2], [Jel3] and [Jel4].

Acknowledgements. The author is grateful to Professor Y. Prokhorov from Steklov Institute of Mathematics for helpful discussions.

2. Terminology.

We assume that ground field $\mathbb{K}$ is algebraically closed of characteristic zero. For an algebraic variety X (variety is here always irreducible) we denote by $Aut(X)$ the group of all regular automorphisms of X and by $Bir(X)$ the group of all birational transformations of X. By $Aut_1(X)$ we mean the group of all birational transformation which are regular in codimension one, i.e., which are regular isomorphisms outside subsets of codimension at least two. If $X \subset \mathbb{P}^n(\mathbb{K})$ then we put $Lin(X) = \{f \in Aut(X) : f = res_X T, T \in Aut(\mathbb{P}^n(\mathbb{K}))\}$. Of course, the group $Lin(X)$ is always an affine group.

Let $f : X - \to Y$ be a rational mapping between projective normal varieties. Then f is determined outside some closed subset F of codimension at least two. If $S \subset X$ and

Date: March 27, 2014.
1991 *Mathematics Subject Classification.* 14 R 10.
Key words and phrases. affine variety, group of automorphisms .
The author was partially supported by the grant of NCN, 2014-2017.

$S \not\subset F$ then by $f(S)$ we mean the set $f(X \setminus F)$. Similarly for $R \subset Y$ we will denote the set $\{x \in X \setminus F : f(x) \in R\}$ by $f^{-1}(R)$.

If $f : X- \to Y$ is a birational mapping and the mapping f^{-1} does not contract any divisor, we say that f is a birational contraction.

An algebraic irreducible variety X of dimension $n > 0$ is called uniruled if there exists an irreducible variety W of dimension $n-1$ and a rational dominant mapping $\phi : W \times \mathbb{P}^1(\mathbb{K}) - \to X$. Equivalently: an algebraic variety X is uniruled if and only if for a generic point in X there exists a rational curve in X through this point.

We say that divisor D is $\mathbb{Q}$-Cartier if for some non-zero integer $m \in \mathbb{Z}$ the divisor mD is Cartier. If every divisor on X is $\mathbb{Q}$-Cartier then we say that X is $\mathbb{Q}$-factorial.

In this paper we treat a hypersurface $H = \bigcup_{i=1}^r H_i \subset X$ as a reduced divisor $\sum_{i=1}^r H_i$, and conversly a reduced divisor will be treated as a hyperserface.

3. Weil divisors on a normal variety.

In this section we recall (with suitable modifications) some basic results about divisors on a normal variety (see e.g., [Iit1]).

Definition 3.1. *Let X be a normal projective variety. We will denote by $Div(X)$ the group of all Weil divisors on X. For $D \in Div(X)$ the set of all Weil divisors linearly equivalent to D, is called a complete linear system given by D and it will be denoted by $|D|$. Moreover we set $L(D) := \{f \in \mathbb{K}(X) : f = 0$ or $D + (f) \geq 0\}$.*

We have the following (e.g., [Iit1], 2.16, p.126)

Proposition 3.2. *If D is an effective divisor on a normal complete variety X then $L(D)$ is a finite dimensional vector space (over $\mathbb{K}$).*

Remark 3.3. The set $|D|$ (if non-empty) has a natural structure of a projective space of dimension $\dim L(D) - 1$. By a basis of $|D|$ we mean any subset $\{D_0, ..., D_n\} \subset |D|$ such that $D_i = D + (\phi_i)$ and $\{\phi_0, ..., \phi_n\}$ is a basis of $L(D)$.

Let us recall the next

Definition 3.4. *If D is an effective Weil divisor on a normal complete variety X then by a canonical mapping given by $|D|$ and a basis ϕ, we mean the mapping $i_{(D,\phi)} = (\phi_0 : ... : \phi_n) : X \to \mathbb{P}^n(\mathbb{K})$, where $\phi = \{\phi_0, ..., \phi_n\} \subset L(D)$ is a basis of $L(D)$.*

Let X be a normal variety and Z be a closed subvariety of X. Put $X' = X \setminus Z$. We would like to compare the group $Div(X)$ and $Div(X')$. It can be easily checked that the following proposition is true (compare [Har], 6.5., p. 133):

Proposition 3.5. *Let $j_{X'} : Div(X) \ni \sum_{i=1}^r n_i D_i \to \sum_{i=1}^r n_i (D_i \cap X') \in Div(X')$. Then $j_{X'}$ is an epimorphism. Moreover it preserves the linear equivalence. If additionally codim $Z \geq 2$ then $j_{X'}$ is an isomorphism.*

Now we define the pull-back of a divisor under a rational map $f : X- \to Y$. Recall that a Cartier divisor can be given by a system $\{U_\alpha, \phi_\alpha\}$ where $\{U_\alpha\}$ is a some open covering of X, $\phi_\alpha \in \mathbb{K}(U_\alpha)$ and $\phi_\alpha/\phi_\beta \in \mathcal{O}^*(U_\alpha \cap U_\beta)$.

Definition 3.6. *Let $f : X \to Y$ be a dominant morphism between normal varieties. Let D be a Cartier divisor on Y given by a system $\{U_\alpha, \phi_\alpha\}$. By the pullback of the divisor D by f we mean the divisor f^*D given by a system $\{f^{-1}(U_\alpha), \phi_\alpha \circ f\}$. More generally if f is a rational map and X_f denotes the domain of f we put $f^*(D) := {j_{X_f}}^{-1}(res_{X_f} f)^* D$.*

Finally let f be as above and D be an arbitrary Weil divisor on Y. Let us assume additionally that $\operatorname{codim} f^{-1}(Sing(Y)) \geq 2$. *Then we have a regular map $f : X_f \setminus W \to Y_{reg}$ (where $W := f^{-1}(Sing(Y))$ and we put $f^*D := {j_{X_f \setminus W}}^{-1} f^*(j_{Y_{reg}}(D))$.*

By a simple verification we have

Proposition 3.7. *Let $f : X - \to Y$ be a dominant rational mapping between complete normal varieties, such that $f^{-1}(Sing(Y))$ has codimension at least two. Then $f^* : Div(Y) \ni D \to f^*D \in Div(X)$ is a well-defined homomorphism, which preserves the linear equivalence. Moreover, $Supp(f^*(D)) = \overline{f^{-1}(Supp(D))}$.*□

Corollary 3.8. *Let f be as above. Let us assume additionally that f is an isomorphisms in codimension one. Then $f^* : Div(Y) \to Div(X)$ is an isomorphisms which preserves the linear equivalence.* □

Finally we have the following important result:

Proposition 3.9. *Let X be a normal complete variety and $f \in Aut_1(X)$. Let D be an effective divisor on X and $f^*D' = D$. Then* $\dim|D| = \dim\ |D'| := n$ *and there exists a unique automorphism $T(f) \in Aut(\mathbb{P}^n(\mathbb{K}))$ such that the folowing diagram commutes*



Proof. First of all let us note that $T(f)$ if exists it is unique. Further, by Corollary 3.8 we have $f^*(|D'|) = |D|$ and f^* transforms any basis of $|D'|$ onto a basis of $|D|$. Let ϕ and ψ be suitable bases such that $i_D = i_{(D,\phi)}$ and $i_{D'} = i_{(D',\psi)}$.

We have $i_{D'} \circ f = (\psi_0, ..., \psi_n) \circ f$. But $f^*(D' + (\psi_i)) = f^*(D') + f^*(\psi_i) = D + (\psi_i \circ f)$. It means that rational functions $(\psi_i \circ f), i = 0, ..., n$ are the basis of $L(D)$. Hence there exists a non-singular matrix $[a_{i,j}]$ such that $\psi_i \circ f = \sum_{j=0}^{n} a_{i,j}\phi_j$. Now it is clear that it is enough to take as $T(f)$ the projective automorphism of $\mathbb{P}^n(\mathbb{K})$ given by the matrix $[a_{i,j}]$. □

Corollary 3.10. *Let G be a subgroup of $Aut_1(X)$ such that $G^*D = D$ for some effective divisor D. Let us denote $\overline{i_D(X)} = X' \subset \mathbb{P}^n(\mathbb{K})$, $n = \dim\ |D|$. Then there is a natural homomorphism $T : G \to Lin(X')$. Moreover, if D is very big (i.e., the mapping i_D is a birational embedding) , then T is a monomorphism.*

Proof. It is enough to take above $D' = D$ and $\phi = \psi$. The last statement is obvious. □

Remark 3.11. In our application we deal only with normal $\mathbb{Q}-$factorial varieties. Hence we can restrict here only to $\mathbb{Q}-$Cartier divisors. However the author thinks that the language of Weil divisors is more natural here.

4. Varieties with good covers.

We begin this section by recalling the definition of a big divisor (see [K-M], p. 67):

Definition 4.1. *Let X be a projective n-dimensional variety and D a Cartier divisor on X. The divisor D is called big if* $\dim H^0(X, \mathcal{O}_X(kD)) > ck^n$ *for some $c > 0$ and $k >> 1$.*

If $f : X \to Y$ is a birational morphisms and D is a big (Cartier) divisor, then also its pullback $f^*(D)$ is big. Indeed the line bundle $\mathcal{O}_X(mf^*(D)) = f^*\mathcal{O}_Y(mD)$ has at least as many sections as the bundle $\mathcal{O}_Y(mD)$. We show later that it is also true for suitable birational mappings (see Lemma 4.5). We have a following characterization of big divisors (see [K-M], Lemma 2.60, p. 67):

Proposition 4.2. *Let X be a projective n-dimensional variety and D a Cartier divisor on X. Then the following are equivalent:*

1) D is big,

2) $mD \sim A + E$, where A is ample and E is effective Cartier divisor,

3) for $m >> 0$ the rational map ι_{mD} associated with the system $|mD|$ is a birational embedding,

4) the image of ι_{mD} has dimension n for $m >> 0$.

In the sequel we need the following observation:

Lemma 4.3. *Let X be a smooth projective variety and let $D = \sum_{i=1}^r a_i D_i$ be a big divisor on X. Then $Supp(D) = \sum_{i=1}^r D_i$ is also a big divisor on X.*

Proof. Let $a = max_{i=1,...,r}\{a_i\}$ and $b_i = a - a_i$. The divisor $E = \sum_{i=1}^r b_i D_i$ is effective. By Proposition 4.3, p. 2) the divisor $D + E = aSupp(D)$ is also big. Hence we conclude by 3) of Proposition 4.3. □

Definition 4.4. *Let X be a normal projective variety and let D be a Weil divisor on X. We say that X is very big if the rational map ι_D associated with the system $|D|$ is a birational embedding. We say that D is big if for some $m \in \mathbb{N}$ the divisor mD is very big.*

We have the following simple lemma:

Lemma 4.5. *Let X, Y be normal projective varieties and let D be an effective big divisor on Y. Let $\phi : X - \to Y$ be a birational mapping such that* $\operatorname{codim} \phi^{-1}(Sing(Y)) \geq 2$. *Then the divisor $\phi^*(D)$ is also big.*

Proof. It is enough to assume that D is very big and prove that then $\phi^*(D)$ is also very big. Take $f_0 = 1$ and let divisors $\{D + (f_0), D + (f_1), ..., D + (f_s)\}$, where $f_i \in \mathbb{K}(Y), i = 0, ..., s$ form a basis of a system $|D|$. By the assumption the regular mapping $\Psi : Y \setminus Supp(D) \ni x \mapsto (f_1(x), ..., f_s(x)) \in \mathbb{K}^s$ is a birational morphism. The system $|\phi^*(D)|$ contains divisors $\{\phi^*(D), \phi^*(D) + (f_1 \circ \phi), ..., \phi^*(D) + (f_s \circ \phi)\}$. Since rational functions $1, f_1 \circ \phi, ..., f_s \circ \phi$ are linearly independent, we can extend them to some basis B of $L(\phi^*(D))$. Let $\Psi' : X \setminus |Supp(\phi^*(D))| \to \mathbb{K}^N$ be a mapping given by a system $|\phi^*(D)|$ and the basis B. The mapping Ψ' after composition with a suitable projection $\mathbb{K}^N \to \mathbb{K}^s$, is equal to $\Psi \circ \phi$. Since the latter mapping is birational, the mapping Ψ' is also birational. □

We shall use:

Definition 4.6. *Let X be an (open) variety. We say that X has a good cover Y, if there exists a completion $\overline{X}$ of X and a smooth projective variety Y with a birational morphism $g : Y \to \overline{X}$ such that:*

1) $D := g^{-1}(\overline{X} \setminus X)$ *is a big hypersurface in* Y*,*

2) $Aut(X) \subset Aut(Y \setminus D)$*, i.e., every automorphism of* X *can be lifted to an automorphism of* $Y \setminus D$*.*

Our next aim is to show that quasi-affine varieties have good covers.

Proposition 4.7. *Let* X *be a quasi-affine variety.Then* X *has a good cover.*

Proof. By the assumption there is an affine variety X_1 such that $X \subset X_1$ is an open dense subset. Since X_1 is affine, we can assume that it is a closed subvariety of some $\mathbb{K}^N$. Denote by $\overline{X}$ the projective closure of X_1 in $\mathbb{P}^N$. Let π_∞ be the hyperplane at infinity in $\mathbb{P}^N$ and $V := \overline{X}.\pi_\infty$ be a divisor at infinity on $\overline{X}$. Of course V is a big (even very ample) Cartier divisor.

Let $h : Y \to \overline{X}$ be a canonical desingularization of $\overline{X}$ (see e.g., [Kol], [Włο]). Then $h_{|h-1(X)} : h^{-1}(X) \to X$ is a canonical desingularization of X. In particular every automorphism of X has a lift to an automorphism of $h^{-1}(X)$, i.e., $Aut(X) \subset Aut(h^{-1}(X)) = Aut(Y \setminus h^{-1}(S))$. Since V is a big divisor, so is its pullback $h^*(V)$.

Let $Z := Y \setminus h^{-1}(X)$. It is a closed subvariety of Y. Let J_Z be the ideal sheaf of Z and let $f : Y' \to Y$ be a canonical principalization of J_Z (see e.g., [Kol], [Włο]). Thus $D := f^{-1}(Z)$ is a hypersurface, which contains a big hypersurface $V' = Supp(f^*h^*(V))$. Since $D = V' + E$, where E is effective divisor, the hypersurface D is also big by Lemma 4.3.

Finally if we take $g = f \circ h : Y' \to \overline{X}$, then conditions 1) and 2) of Definition 4.6 are satisfied. □

5. The Quasi Minimal Model

In this section following [P-S], we introduce the notion of quasi-minimal models (for details see [P-S]). This is a weaker analog of a usual notion of minimal models which has an advantage that to prove its existence we do not need the full strength of the Minimal Model Program.

Definition 5.1. *(see* [P-S]*) An effective* $\mathbb{Q}$*-divisor* M *on a variety* X *is said to be* Q*-movable if for some* $n > 0$ *the divisor* nM *is integral and generates a linear system without fixed components. Let* X *be a projective variety with* $\mathbb{Q}$*-factorial terminal singularities. We say that* X *is a quasi-minimal model if there exists a sequence of* $\mathbb{Q}$*-movable* $\mathbb{Q}$*-divisors* Mj *whose limit in the Neron-Severi space* $NSW_{\mathbb{Q}}(X) = NSW(X) \otimes \mathbb{Q}$ *is* K_X*.*

By the recent progress in the minimal model program (see [Bir], [BCHK], [P-S]) every non-uniruled smooth variety has a quasi-minimal model. In fact if we ran MMP on X and we do all possible divisorial contractions (and all necessary flips) we achieve a quasi-minimal model Y, moreover we obtain the mapping $\phi : X - \to Y$ which is a composition of divisorial contractions ans flips, in particular it is a birational contraction, i.e., the mapping ϕ^{-1} does not contract any divisor (cf. [P-S], section 4, Corollary 4.5):

Theorem 5.2. *Let* X *be a smooth projective non-uniruled variety. Then there is a quasi-minimal model* Y *and a birational contraction* $\phi : X - \to Y$*.*□

Quasi minimal models have the following very important property (cf. [P-S], section 4, Proposition 4.6):

Theorem 5.3. *Let* X *be a quasi-minimal model. Then* $Bir(X) = Aut_1(X)$*.*□

6. Main Result.

Now we can start our proof. The first step is

Proposition 6.1. *Let X be a normal complete non-uniruled variety and H be a big hypersurface on X. Then the group $Stab_X(H) = \{f \in Aut_1(X) : f^*H = H\}$ is finite.*

Proof. For some $m \in \mathbb{N}$ the divisor mD is very big. We have $Stab_X(H) = \{f \in Aut_1(X) : f^*H = H\} = Stab_X(mH) = \{f \in Aut_1(X) : f^*(mH) = mH\}$.

By the assumption the variety $X' = \overline{i_{mH}(X)}$ is birationally equivalent to X. In view of Corollary 3.10 it is enough to prove that the group $Lin(X')$ is finite. Since X is non-uniruled, the X' is non-uniruled, too. But the group $Lin(X')$ is an affine group and if it is infinite, then by Rosenlicht Theorem (see [Ros]) we have that X' is ruled - which is impossible. □

Now we can prove our main result:

Theorem 6.2. *Let X be an open variety with a good cover. If the group $Aut(X)$ is infinite, then X is uniruled.*

Proof. Assume that $Aut(X)$ is infinite. Let $f : \overline{Y} \to \overline{X}$ be a good cover of X and take $Y = f^{-1}(X)$. Then $Aut(Y)$ is also infinite. We have to prove that X is uniruled. To do this it suffices to prove that Y is uniruled.

Assume that Y is not uniruled. By Theorem 5.3 there exists a quasi-minimal model Z and a birational contraction $\phi : \overline{Y} - \to Z$. Take $\psi = \phi^{-1}$. The mapping ψ is a regular mapping outside some closed subset F of codimension ≥ 2. By the Zariski Main Theorem the mapping ψ restricted to $Z \setminus F$ is an embedding.

Take a mapping $G \in Aut(Y)$, in fact $G \in Bir(\overline{Y})$. The mapping G induces a birational mapping $g \in Bir(Z)$. Since $Bir(Z) = Aut_1(Z)$ we have $g \in Aut_1(Z)$. The mapping g is a morphisms outside a closed subset R of codimension ≥ 2, moreover sets $\overline{g^{-1}(R)} = R_1$ and $\overline{g^{-1}(F)} = F_1$ have also codimension at least two (g does not contract divisors).

The mapping ψ embeds the set $V := Z \setminus (F \cup F_1 \cup R \cup R_1)$ into Y, denote $U := \psi(Z \setminus (F \cup F_1 \cup R \cup R_1))$. Under this identification the mapping g restricted to V corresponds to the mapping G restricted to U.

Let $D = \overline{Y} \setminus Y$ be a big hypersurface, as in the definition of a good cover. The hypersurface $D' := \psi^*(D)$ is also big (see Lemma 4.5) and $D' \cap V$ corresponds to $D \cap U$. Since $G(U \setminus D) = U \setminus D$ we have that g transforms irreducible components of $D' \cap V$ onto irreducible components of $D' \cap V$. In particular $g^*(D') = D'$. This means that $Aut(Y) \subset Stab_{D'}(Z) \subset Aut_1(Z)$. By Proposition 6.1 this contradicts our assumption. □

Corollary 6.3. *Let X be a quasi-affine (in particular affine) variety. If the group $Aut(X)$ is infinite, then X is uniruled.*

References

[Bir] Birkar, C, *Existence of log canonical flips and a special LMMP*, Publ. Math. Inst. Hautes Etudes Sci., 325368, (2012).

[BCHK] Birkar, C; Cascini, P; Hacon, C.; McKernan, J, *Existence of minimal models for varieties of log general type*, Journal of the AMS, 23 (2): 405468, (2010).

[Har] Hartshorne, R, *Algebraic Geometry*, Springer Verlag, New York, 1987.

[Iit1] Iitaka, S, *Algebraic Geometry*, Springer Verlag, New York, 1982.

[Iit2] Iitaka, S, *On logarithmic Kodaira dimension of algebraic varieties.* In: Complex Analysis and Algebraic Geometry, 178-189, Tokyo: Iwanami 1977.

[Jel1] Jelonek, Z, Irreducible Identity Sets for Polynomial Automorphisms, *Math.Z.* **212**, 601-617 (1993).
[Jel2] Jelonek, Z, Affine smooth varieties with finite group of automorphisms, *Math. Z.* **216** , 575-591 (1994).
[Jel3] Jelonek, Z, The group of automorphisms of an affine non-uniruled surface, **Univ. Iaegel. Acta Math. 32**, 65-68 (1995),
[Jel4] Jelonek, Z, The group of automorphisms of an affine non-uniruled variety, *Seminari di Geometria*, Uniwersita degli Studi di Bologna, 169-180, (1996)
[K-M] Kollar, J, Mori, S, Birational geometry of algebraic varieties, Cambridge University Press, 1998.
[Kol] Kollar, J, Lectures on Resolution of Singularities, Annals of Mathematics Studies 166, (2007).
[P-S] Prokhorov, Y, Shramov, C, Jordan property for groups of birational selfmaps, arXiv: 1307.1784v2 (2013).
[Ros] Rosenlicht, M, Some Basic Theorems on Algebraic Groups. *American Journal of Mathematics*, **78**, 401-443, (1956).
[Wło] Włodarczyk, J, Simple Hironaka resolution in characteristic zero, *J. Amer. Math. Soc.* 18 , 779-822, (2005).

(Z. Jelonek) Instytut Matematyczny, Polska Akademia Nauk, Śniadeckich 8, 00-956 Warszawa, Poland

E-mail address: najelone@cyf-kr.edu.pl